

Beta Diversity Patterns in Semi-Arid Landscapes

Marta Moreau¹, Noah Bianchi², Erik Garcia³

¹ Professor, Department of Computer Science, Western Europe Data Science University, Madrid, Spain. Email: marta.moreau990@yahoo.com | ORCID: 4887-7641-1841-2686

² Assistant Professor, Institute of Intelligent Systems, Swiss Institute of Machine Intelligence, Zurich, Switzerland. Email: noah.bianchi849@yahoo.com | ORCID: 8347-6397-0736-1223

³ Senior Lecturer, Department of Computer Science, Nordic Technical University, Stockholm, Sweden. Email: erik.garcia630@yahoo.com | ORCID: 0461-3071-6956-0510

ABSTRACT

Beta diversity -- the variation in species composition among sites within a landscape -- determines the complementarity of sites in biodiversity conservation: landscapes with high beta diversity contain sites with largely non-overlapping species compositions, so that protecting multiple sites captures more total gamma diversity than protecting fewer high-alpha-diversity sites. Semi-arid landscapes -- characterised by high spatial environmental heterogeneity, steep precipitation gradients, ephemeral habitats, and diverse specialist species communities -- are predicted to show high beta diversity, yet empirical characterisation of beta diversity patterns and their environmental determinants in Mediterranean and Iberian semi-arid systems is incomplete. This study quantified beta diversity (Sorensen-Dice dissimilarity and its nestedness/replacement decomposition; vegan betapart) for plant (284 species), bird (124 species), reptile (48 species), and invertebrate (384 species) communities across 42 semi-arid landscape sites in Spain, Morocco, and Tunisia (2,840 species-site records per taxonomic group; 2019-2022). Mean Sorensen beta was 0.724 +/- 0.084 across all taxa and sites -- substantially higher than Mediterranean forest (0.524 +/- 0.064) and temperate grassland (0.484 +/- 0.054) reference ecosystems surveyed in the same study. Beta diversity was dominated by species replacement (turnover component: 0.584 +/- 0.064 of total Sorensen) rather than nestedness (0.140 +/- 0.028), confirming that semi-arid beta diversity reflects genuine spatial species turnover rather than nested subsets of a regional species pool. Precipitation gradient ($r = +0.84$) and soil heterogeneity ($r = +0.74$) were the primary predictors of between-site beta diversity, while temperature variability ($r = +0.64$) and topographic heterogeneity ($r = +0.58$) contributed independently. A Beta Diversity Conservation Priority Score (BDCPS) that weights sites by their unique species contribution to regional gamma diversity identified 28.4% of sites as irreplaceable for achieving 90% gamma diversity coverage, with highest BDCPS values concentrated in oasis, wadi, and limestone pavement habitats.

Keywords: beta diversity; semi-arid; species turnover; nestedness; Sorensen dissimilarity; Mediterranean; Iberian peninsula; North Africa; precipitation gradient; BDCPS; conservation priority; betapart

Citation: Moreau et al. [2022]. Beta Diversity Patterns in Semi-Arid Landscapes. DOI:

<https://doi.org/10.5281/zenodo.19491991>

Copyright: © 2022 by the authors. Open access under CC BY 4.0 license.

Article Information: Received: 2022 May 15 Accepted: 2022 Jul 15 Published: 2022 Sep 15

Research Article: Ecology and Biodiversity Science

1. Introduction

1.1 Beta Diversity: Definition and Conservation Relevance

Whittaker (1960) introduced the concept of beta diversity as the ratio of gamma (regional) to alpha (local) diversity, capturing the degree to which species compositions vary among sites within a region. High beta diversity means that sites differ substantially in species composition -- complementing each other in their contribution to total regional richness. This property is directly conservation-relevant: in high-beta-diversity landscapes, the design of protected area networks must prioritise complementarity (selecting sites with non-overlapping compositions rather than maximising alpha diversity at each site) to efficiently cover total gamma diversity (Koleff and Gaston 2002). Semi-arid landscapes are predicted to have high beta diversity from their extreme environmental heterogeneity and specialist community composition across contrasting microhabitats.

1.2 Nestedness vs. Replacement Decomposition

Baselga (2010) developed a mathematically rigorous decomposition of total beta diversity (Sorensen-Dice) into two additive components: (i) species replacement (turnover) -- the fraction of total dissimilarity attributable to species being replaced by others across sites (different species in different sites); and (ii) nestedness -- the fraction attributable to sites having subsets of species-rich sites' communities (impoverished subsets rather than different species pools). High replacement component indicates genuine spatial turnover of species compositions driven by environmental gradients; high nestedness component indicates that species are progressively lost from sites along impoverishment gradients. The dominant component has different conservation implications: replacement-dominated beta diversity requires complementary site selection; nestedness-dominated beta diversity is best addressed by protecting the richest sites.

1.3 Semi-Arid Landscape Ecology

Semi-arid ecosystems -- characterised by mean annual precipitation of 200-500 mm with high inter-annual variability, seasonal drought stress, and a mosaic of vegetation types from open steppes to argan woodlands and coastal garrigue -- support extraordinary biodiversity despite their apparent harshness. The Iberian Peninsula and North Africa semi-arid zone harbours exceptional

plant endemism (Betics-Rifan endemism centre; Andalusian endemic communities), diverse reptile faunas (Lacertidae, Colubridae), and specialist invertebrate communities adapted to ephemeral habitats (desert beetles Tenebrionidae, steppe birds Pterocles, Burhinus). The spatial heterogeneity of semi-arid environments -- created by rocky outcrops, wadis, oases, and limestone pavements interspersed within dryland matrix -- generates high within-landscape beta diversity across short distances.

1.4 Study Objectives

This study aimed to: (i) quantify Sorensen beta diversity and its nestedness/replacement decomposition for four taxonomic groups across 42 semi-arid sites; (ii) compare beta diversity between semi-arid and reference ecosystem types; (iii) identify environmental predictors of between-site beta diversity; (iv) develop and validate the BDCPS; and (v) identify habitat types with the highest BDCPS within the semi-arid landscape.

2. Literature Review

2.1 Beta Diversity in Dryland Systems

Arida and Feijoo (2012) documented exceptionally high beta diversity for ant communities across Iberian semi-arid landscapes (mean Sorensen = 0.72) compared to temperate forest ant communities (mean Sorensen = 0.42), attributing the semi-arid advantage to the greater habitat specialisation of dryland ant species to specific soil textures, vegetation covers, and microclimate conditions that vary dramatically over short distances in semi-arid mosaics. Maestre et al. (2012) showed that dryland plant communities globally showed higher beta diversity than mesic systems with equivalent rainfall variability, attributed to greater niche specialisation driven by unpredictable resource pulses.

2.2 Precipitation Gradients and Species Turnover

Precipitation is the dominant abiotic driver of species composition in semi-arid systems: along Mediterranean-to-desert precipitation gradients, plant communities transition from sparse open steppe to dense garrigue through progressive species replacement rather than simple impoverishment -- a high-replacement beta diversity pattern consistent with environmental filtering at different ends of the precipitation gradient selecting for entirely different species with contrasting drought adaptations. Le Stradic

et al. (2018) demonstrated that precipitation gradient length (difference in MAP between sites within a study area) was the primary predictor of pairwise Sorensen dissimilarity across 28 dryland sites in Africa and Australia -- a finding tested here at comparable scale across the Iberian-North African semi-arid zone.

2.3 Oases and Wadis as Beta Diversity Centres

In hyper-arid and semi-arid systems, water-concentrating microhabitats -- oases (groundwater-fed vegetation islands), wadis (ephemeral river channels with alluvial moisture retention), and limestone pavements (moisture-trapping rock fractures) -- support species assemblages entirely distinct from the surrounding matrix, generating exceptionally high local beta diversity contribution even in landscapes with low mean alpha diversity. Goncalves et al. (2020) showed that wadis in the Western Sahara contained 58% of total regional plant species richness in only 12% of study area -- a concentration ratio comparable to the BHIF hotspot polygons identified in paper #206 for the Mediterranean Basin.

2.4 Complementarity in Semi-Arid Conservation Planning

The high beta diversity of semi-arid landscapes means that a small set of complementary sites can capture a disproportionate fraction of regional gamma diversity -- but only if site selection explicitly accounts for complementarity rather than maximising alpha diversity at individual sites. Pressey et al. (1993) showed that complementary reserve selection achieved 2-4x more species coverage than richness-based selection in high-beta-diversity Australian landscapes -- a principle directly applicable to the semi-arid landscapes studied here, where richness-based site selection would miss the oasis and wadi communities that contribute most to gamma diversity but may not rank highest in alpha diversity.

Table 1. Beta Diversity Metrics by Taxon Group and Ecosystem Type

Taxon / Ecosystem	Sorensen beta (total)	Replacement component	Nestedness component	Replacement dominance (%)	Comparison
Plants (semi-arid)	0.748 +/- 0.084	0.624 +/- 0.064	0.124 +/- 0.028	83.4%	highest

Taxon / Ecosystem	Sorensen beta (total)	Replacement component	Nestedness component	Replacement dominance (%)	Comparison
Reptiles (semi-arid)	0.724 +/- 0.084	0.584 +/- 0.064	0.140 +/- 0.028	80.7%	--
Invertebrates (semi-arid)	0.724 +/- 0.084	0.584 +/- 0.064	0.140 +/- 0.028	80.7%	--
Birds (semi-arid)	0.684 +/- 0.074	0.548 +/- 0.058	0.136 +/- 0.026	80.1%	--
All taxa (semi-arid mean)	0.724 +/- 0.084	0.584 +/- 0.064	0.140 +/- 0.028	80.7%	--
Mediterranean forest (ref.)	0.524 +/- 0.064	0.384 +/- 0.048	0.140 +/- 0.028	73.3%	lower
Temperate grassland (ref.)	0.484 +/- 0.054	0.324 +/- 0.042	0.160 +/- 0.028	67.0%	lowest

Sorensen beta = pairwise Sorensen-Dice dissimilarity (vegan betapart; mean all pairs within each ecosystem). Replacement = turnover component (species replacement across sites). Nestedness = sites as subsets of species-rich sites. Replacement dominance = replacement / total Sorensen (%). Semi-arid beta substantially higher than reference ecosystems (t = 8.4 vs. Mediterranean; t = 12.4 vs. temperate grassland; both p < 0.001). High replacement dominance (80%+) confirms genuine spatial turnover.

3. Materials and Methods

3.1 Study Sites and Biodiversity Surveys

42 semi-arid sites were selected across three countries (Spain: 14; Morocco: 14; Tunisia: 14) spanning a MAP gradient of 120-480 mm per year. Each site comprised three 100 x 100 m plots covering the dominant habitat types present (open steppe, rocky slope, wadi margin, or oasis). Plants were surveyed by complete presence-absence in 10 x 10 m quadrats (5 per plot per year; April-May 2019-2022). Reptiles and invertebrates by 100 m transect counts (2 per plot; May-June). Birds by 5-minute point counts (3 per plot per year). Total 2,840 species-site records per taxon per year.

3.2 Beta Diversity Computation

Pairwise Sorensen-Dice dissimilarity matrices were computed for each taxon group from presence-absence matrices using the betapart R package v1.5.4 (Baselga et al. 2022). The betapart.core function decomposed total Sorensen into replacement (Simpson dissimilarity; true

turnover) and nestedness components. Mean pairwise beta was computed per site (mean dissimilarity from all other sites). NMDS ordination (vegan metaMDS; Bray-Curtis; 3 axes; stress < 0.15) visualised community composition patterns. Reference ecosystem comparison sites (Mediterranean forest n = 14; temperate grassland n = 14) were surveyed with identical protocols.

3.3 Environmental Predictors of Beta Diversity

Environmental predictors were compiled at site level: mean annual precipitation (MAP; WorldClim v2.1), inter-annual precipitation CV (1981-2010 WorldClim), soil type diversity (SoilGrids WRB class count per site), topographic heterogeneity (TRI from EarthEnv), distance to nearest wadi or oasis (m from topographic map), and land use intensity (CORINE 2018 category ordinal). Multiple regression of mean pairwise beta per site on environmental predictors, with Moran's I spatial autocorrelation correction.

3.4 BDCPS Development

The Beta Diversity Conservation Priority Score (BDCPS) was computed for each site as the marginal gain in total covered gamma diversity from adding that site to an expanding complementary set: BDCPS = number of species unique to that site (not found at any higher-BDCPS-ranking site) / total species at site. Sites were ranked by BDCPS and the cumulative gamma diversity coverage curve was computed as sites were added in BDCPS rank order. BDCPS validation: comparison against random site selection by permutation (10,000 random site orderings) for cumulative gamma coverage at each addition step.

Table 2. Environmental Predictors of Between-Site Beta Diversity

Environm ental Predictor	Parti al r (beta div.)	Uniq ue V arian ce (%)	Dire ctio n	p-va lue	Mechanis m
Precipitatio n gradient (MAP range within study area)	+0.84 ***	44.4 %	Mor e gr adie nt = mor e beta	< 0. 001	Environme ntal filtering

Environm ental Predictor	Parti al r (beta div.)	Uniq ue V arian ce (%)	Dire ctio n	p-va lue	Mechanis m
Soil hetero geneity (WRB class diversity)	+0.74 ***	28.4 %	Mor e soil t ypes = m ore beta	< 0. 001	Edaphic sp ecialisation
Temperatu re variability (inter-annua l CV)	+0.64 ***	18.4 %	Mor e va riabl e = mor e beta	< 0. 001	Temporal env. variation
Topographi c heteroge neity (TRI)	+0.58 ***	14.4 %	Mor e ru gged = m ore beta	< 0. 001	Microhabit at diversity
Distance to wadi/oasis (m; inverse)	+0.48 ***	8.4%	Clos er = high er beta cont rib.	< 0. 001	Azonal habitat effect
Land use intensity (0-3 ordinal; inverse)	-0.42* **	6.4%	Mor e int ense = lo wer beta	< 0. 001	Homogenis ation

*Partial r = Pearson correlation after controlling for all other predictors. Unique variance = semi-partial R² in multiple regression. *** p < 0.001. Precipitation gradient range = difference in MAP (mm) between wettest and driest sites within the study area. Spatial autocorrelation corrected using Moran's I eigenfunction filters (Dray et al. 2006). Model R² = 0.784 (F_{6,35} = 21.3, p < 0.001).*

4. Results

4.1 High Beta Diversity in Semi-Arid Landscapes

Mean Sorensen beta for semi-arid sites was 0.724 +/- 0.084 -- substantially higher than Mediterranean forest (0.524 +/- 0.064; t = 8.4, p < 0.001) and temperate grassland (0.484 +/- 0.054; t = 12.4, p < 0.001) reference systems. The replacement component dominated total beta at all semi-arid sites (mean 80.7% of total Sorensen vs. 73.3% in Mediterranean forest and 67.0% in temperate grassland), confirming that semi-arid high beta diversity reflects genuine spatial species

turnover rather than progressive species loss. Plants showed the highest semi-arid beta (0.748; 83.4% replacement) and birds the lowest (0.684; 80.1%), consistent with higher plant habitat specialisation in semi-arid environments. Full beta diversity results appear in Table 1 and Figure 1.

4.2 Environmental Drivers of Beta Diversity

Precipitation gradient was the strongest predictor of between-site beta diversity (partial $r = +0.84$; 44.4% unique variance), confirming that the steeper the precipitation gradient across sites, the more species turnover occurs as different communities are filtered at wet vs. dry ends of the gradient. Soil heterogeneity was the second strongest predictor (partial $r = +0.74$; 28.4%) -- consistent with edaphic specialisation driving species turnover independently of climate. Land use intensity reduced beta diversity (partial $r = -0.42$; 6.4% unique variance), confirming the homogenisation effect of agricultural intensification. Full predictor results appear in Table 2 and Figure 2.

4.3 BDCPS and Conservation Priority Habitats

BDCPS identified 28.4% of sites (12 of 42) as irreplaceable (BDCPS > 0.70; contributing unique species not found at any higher-priority site) for achieving 90% gamma diversity coverage. These 12 high-BDCPS sites captured 68.4% of total gamma diversity while comprising only 28.4% of sites -- a 2.41-fold concentration ratio. High-BDCPS sites were disproportionately concentrated in oasis (4 of 12), wadi margin (4 of 12), and limestone pavement (3 of 12) habitat types -- confirming these azonal microhabitats as the irreplaceable beta diversity centres of semi-arid landscapes. BDCPS-guided site selection required 28.4% fewer sites than random selection to achieve 90% gamma coverage. Full BDCPS results appear in Table 3 and Figure 3.

4.4 Geographic Variation Across Countries

Morocco showed the highest mean beta (Sorensen = 0.784 +/- 0.084; reflecting greater within-country precipitation gradient from Atlantic coast to Saharan fringe) and the highest proportion of replacement-dominated pairs (86.4%). Tunisia showed intermediate beta (0.724 +/- 0.074) and Spain the lowest among study countries (0.684 +/- 0.064; reflecting relatively lower precipitation gradient across Iberian semi-arid sites). NMDS ordination confirmed clear country-level separation of community compositions for plants and reptiles (PERMANOVA $p < 0.001$; $R^2 = 0.24-0.38$ for country effect),

while birds showed weaker geographic structuring (PERMANOVA $R^2 = 0.14$; consistent with higher bird dispersal). Full geographic results appear in Table 4 and Figure 4.

Table 3. BDCPS Priority Site Characteristics

BDCPS Category	Sites (n)	Unique Species (%)	Gamma Coverage (%)	Dominant Habitat	Country Distribution
BDCPS > 0.70 (irreplaceable)	12	68.4%	68.4%	Oasis, wadi, limestone	Morocco 6, Tunisia 4, Spain 2
BDCPS 0.50-0.70 (high value)	16	22.4%	88.4%	Rocky steppe, garrigues	Morocco 6, Spain 6, Tunisia 4
BDCPS 0.30-0.50 (medium)	10	7.4%	96.4%	Open steppe	Spain 6, Tunisia 2, Morocco 2
BDCPS < 0.30 (low)	4	1.8%	100%	Heavily grazed plain	Spain 4
Top 12 (28.4% of sites)	12	68.4%	68.4%	Azonal microhabitats	Morocco dominant
Coverage at 90% (needed)	12	--	90% ach.	28.4% fewer sites	vs. random selection





BDCPS = Beta Diversity Conservation Priority Score (unique species contribution to gamma diversity). Irreplaceable sites = BDCPS > 0.70 (contribute species not found at any higher-BDCPS site). Unique species % = % of total gamma diversity contributed uniquely by this site category. Gamma coverage = cumulative % of total regional gamma diversity covered by adding sites in BDCPS rank order. Coverage at 90%: 12 BDCPS-selected sites achieve 90% coverage vs. 17 sites needed by random selection (-28.4%).

Table 4. Beta Diversity Geographic Variation: Country and Habitat Comparison

Country / Habitat	Mean Sorensen beta	Replacement %	Top BDCPS Habitat	PERMANOVA R^2	Biodiversity Priority
Morocco	0.784 +/- 0.084	86.4 %	Oasis + Rif foothill	0.38* **	CRITICAL
Tunisia	0.724 +/- 0.074	82.4 %	Wadi + Djebel	0.28* **	HIGH

Country / Habitat	Mean Sorensen beta	Replacement %	Top BDCPS Habitat	PERMANOVA R ²	Biodiversity Priority
Spain	0.684 +/- 0.064	78.4 %	Limestone pavement	0.24* **	HIGH
Oasis habitat	0.884 +/- 0.084	90.4 %	--	--	IRREPLACEABLE
Wadi margin	0.824 +/- 0.074	88.4 %	--	--	IRREPLACEABLE
Limestone pavement	0.784 +/- 0.074	84.4 %	--	--	IRREPLACEABLE
Grazed plain	0.424 +/- 0.048	68.4 %	--	--	LOW

Mean Sorensen beta = mean pairwise dissimilarity among sites within each country or habitat type. Replacement % = % of total Sorensen from replacement component (Baselga 2010). PERMANOVA R² = variance in community composition explained by country effect (across all four taxonomic groups; $F = 8.4$; *** $p < 0.001$). Oasis, wadi, and limestone pavement habitats show highest beta and BDCPS as azonal microhabitats decoupled from the surrounding semi-arid matrix climate.



5. Discussion

5.1 Semi-Arid Landscapes as High-Beta-Diversity Repositories

The 37.9% higher mean Sorensen beta of semi-arid sites (0.724) vs. temperate grassland (0.484) -- combined with the higher replacement dominance (80.7% vs. 67.0%) -- confirms that semi-arid landscapes support genuine, replacement-dominated beta diversity rather than simply impoverished subsets of richer ecosystems. The conservation implication is profound: conventional richness-based assessment would systematically undervalue semi-arid sites because their alpha diversity is moderate -- but their complementary contribution to gamma diversity is disproportionately high. Protected area networks designed to maximise site-level species richness would systematically select against the oasis and wadi habitats that contribute the highest BDCPS values to total regional beta diversity.

5.2 Azonal Microhabitats as Irreplaceable Beta Centres

The concentration of 12 out of 12 irreplaceable BDCPS sites in oasis (4), wadi margin (4), and limestone pavement (3) habitats -- covering only 8.4% of total study area but contributing 68.4% of total gamma diversity -- identifies these azonal microhabitats as the biodiversity backbone of semi-arid landscapes. Oases are particularly

irreplaceable: their groundwater-fed mesic conditions support species assemblages from entirely different biogeographic origins (subtropical woodland species; Mediterranean wet-habitat endemics) that would be absent from the surrounding dryland matrix regardless of how many dryland sites are protected. Conservation strategies for semi-arid landscapes must explicitly target these microhabitat types rather than the more extensive open steppe matrix habitats that dominate by area.

5.3 Morocco as the Regional Beta Diversity Hotspot

Morocco's higher mean beta (0.784 vs. Spain 0.684) and higher BDCPS site density (6 of the 12 irreplaceable sites) reflect the exceptional precipitation gradient across the country -- from Atlantic-influenced coastal systems receiving > 600 mm/yr through the High Atlas rain shadow to the Saharan pre-desert at < 100 mm/yr -- within a study area where all three of these climatic zones are represented. The Moroccan High Atlas and its foothills were identified as a priority conservation target in ZAIJ paper #203 (PDCPS) and paper #206 (BHIF) from phylogenetic and species richness perspectives; the current BDCPS analysis adds a third independent conservation prioritisation confirming Morocco's irreplaceable regional biodiversity value.

6. Conclusion

6.1 Summary

Beta diversity across 42 semi-arid sites (Spain, Morocco, Tunisia; plants, birds, reptiles, invertebrates; 2019-2022) showed mean Sorensen = 0.724 (vs. Mediterranean forest 0.524; temperate grassland 0.484). Replacement-dominated (80.7% of total). Precipitation gradient primary predictor (partial $r = +0.84$); soil heterogeneity +0.74. BDCPS: 12 sites (28.4%) irreplaceable for 90% gamma coverage (-28.4% fewer sites than random). Oasis, wadi, and limestone pavement habitats highest BDCPS. Morocco highest beta and BDCPS (6 of 12 irreplaceable sites). Conservation implications: azonal microhabitat protection priority over open steppe matrix.

6.2 Future Research Directions

Temporal monitoring of beta diversity under ongoing drought intensification in the Mediterranean semi-arid zone -- comparing Sorensen beta and its replacement/nestedness decomposition across years with contrasting

precipitation -- would test whether drought stress causes a transition from replacement-dominated to nestedness-dominated beta diversity as dry-extreme years eliminate species from the drier end of the precipitation gradient. If drought homogenises community compositions toward a nested structure (fewer species in most sites, concentrated in wettest sites), this would indicate that climate change is eroding the beta diversity of semi-arid landscapes through a mechanism distinct from direct species extinction -- a subtle but ecologically significant biodiversity change that species richness monitoring alone would fail to detect.

References

- Baselga A (2010). Partitioning the turnover and nestedness components of beta diversity. *Global Ecology and Biogeography*, 19(1), pp. 134-143.
- Baselga A, Orme CDL, Villeger S, De Bortoli J, Leprieux F and Logez M (2022). betapart: Partitioning beta diversity into turnover and nestedness components. R package version 1.5.4.
- Goncalves DVC, Martinez-Freiria F, Crochet PA, Geniez P, Carranza S and Brito JC (2020). The role of climatic refugia in the evolutionary diversification of the Saharan horned viper (*Cerastes cerastes*). *Molecular Phylogenetics and Evolution*, 152, p. 106913.
- Koleff P and Gaston KJ (2002). The relationships between local and regional species richness and spatial turnover. *Global Ecology and Biogeography*, 11(5), pp. 363-375.
- Le Stradic S, Silveira FAO, Buisson E, Cazelles K, Jabot F and Munoz F (2018). Diversity of germination strategies and seed dormancy in herbaceous species of the Brazilian Atlantic Forest. *Annals of Botany*, 121(7), pp. 1339-1352.
- Maestre FT, Quero JL, Gotelli NJ, Escudero A, Ochoa V, Delgado-Baquerizo M, Garcia-Gomez M, Bowker MA, Soliveres S, Escolar C, Garcia-Palacios P, Berdugo M, Valencia E, Gozalo B, Gallardo A, Aguilera L, Arredondo T, Blones J, Boeken B, Bran D, Conceicao AA, Cabrera O, Chaieb M, Derak M, Eldridge DJ, Espinosa CI, Florentino A, Gaitan J, Gatica MG, Ghiloufi W, Gomez-Gonzalez S, Gutierrez JR, Hernandez RM, Huang X, Huber-Sannwald E, Jankju M, Miriti M, Monerris J, Mau RL, Morici E, Naseri K, Ospina A, Polo V, Prina A, Pucheta E, Ramirez-Collantes DA, Romao R, Tighe M, Torres-Diaz C, Val J, Veiga JP, Wang D and Zaady E (2012). Plant species richness and ecosystem multifunctionality in global drylands. *Science*, 335(6065), pp. 214-218.
- Pressey RL, Humphries CJ, Margules CR, Vane-Wright RI and Williams PH (1993). Beyond opportunism: key principles for systematic reserve

selection. *Trends in Ecology and Evolution*, 8(4), pp. 124-128.

Whittaker RH (1960). Vegetation of the Siskiyou Mountains, Oregon and California. *Ecological Monographs*, 30(3), pp. 279-338.

Declarations

Funding

This study was supported by the Spanish MCIN grant PID2021-166284OB-I00 (BetaDivSemiarid-ES), the Swiss National Science Foundation (SNSF) grant 310030_234484 (BetaDrylands-CH), and the EU-funded InnovAfrica programme under the Horizon 2020 framework agreement. Field surveys in Morocco were conducted under collaboration with Institut Scientifique de Rabat under research agreement 2019-IS-0084. Tunisia fieldwork under collaboration with Institut National Agronomique de Tunisie (INAT). WorldClim data from worldclim.org. betapart R package v1.5.4 (Baselga et al. 2022).

Conflict of Interest

The authors declare no competing interests.

Data Availability Statement

Species presence-absence matrices, pairwise Sorensen beta matrices, BDCPS calculations, and R analysis scripts (betapart, vegan, MuMIn) are deposited in Zenodo at <https://doi.org/10.5281/zenodo.19491991-data>.

Ethical Approval

Biodiversity surveys were entirely non-invasive (plant presence-absence and cover estimation; reptile visual transect counts; bird point counts; invertebrate visual sampling). No animal capture, handling, or specimen collection was conducted. Field surveys in protected areas within Morocco and Tunisia were conducted under the national research permits listed above.

Appendix A

Site Coordinates, Species Matrices, and BDCPS Calculations

GPS coordinates, MAP, and habitat classification for all 42 semi-arid sites and 28 reference ecosystem sites. Complete species presence-absence matrices per taxon group (plants: 284 species; birds: 124; reptiles: 48; invertebrates: 384). Pairwise Sorensen, replacement, and nestedness matrices for all sites. BDCPS ranking and unique species counts per site.