

Community Assembly Rules in Island Ecosystems

Eva Kovacs¹, Eva Schmidt², Nina Schmidt³

¹ Professor, Department of Computer Science, Nordic Technical University, Stockholm, Sweden. Email: eva.kovacs563@gmail.com | ORCID: 9669-8638-1621-3156

² Professor, Department of Machine Learning, Baltic AI Research University, Tallinn, Estonia. Email: eva.schmidt735@gmail.com | ORCID: 3100-1777-9911-0332

³ Professor, Department of Machine Learning, Central European Tech University, Vienna, Austria. Email: nina.schmidt494@gmail.com | ORCID: 5621-5737-7431-5955

ABSTRACT

Island ecosystems -- oceanic islands and habitat islands -- provide natural laboratories for studying community assembly processes because their discrete boundaries, controlled immigration history, and often well-documented geological ages enable testing of assembly theory with a precision impossible in mainland systems. MacArthur and Wilson's (1967) equilibrium theory of island biogeography predicts that island species richness is determined by the dynamic balance between immigration and extinction rates, mediated by island area and isolation -- yet the mechanistic processes determining which species colonise and persist remain incompletely understood. This study tested four non-exclusive assembly rules (environmental filtering, competition-driven limiting similarity, neutral stochastic dispersal, and priority effects) across 42 island communities from three island systems (Macaronesian archipelagos: Canary Islands + Azores, $n = 14$; Mediterranean islands: Balearics + Aegean, $n = 14$; Atlantic coastal archipelagos: UK + Ireland smaller islands, $n = 14$) using 2,840 species-site records (plants, birds, reptiles, invertebrates; 2019-2022) and null model analysis (picante; 999 permutations). Environmental filtering dominated assembly in 68.4% of island communities ($NRI > +1.0$; phylogenetic clustering), competition-driven limiting similarity in 18.4% of communities ($NRI < -1.0$; overdispersion), and stochastic dispersal in 13.2% (NRI not significant). Island area and isolation were jointly significant predictors of NRI (larger, less isolated islands showed stronger competition signal; smaller, more isolated islands showed stronger filtering). An Island Community Assembly Index (ICAI) combining NRI , functional trait clustering, and dispersal limitation evidence predicted extinction risk of endemic species with $AUC = 0.864$ and native species conservation status with $r = +0.74$. Under SSP2-4.5, 48.4% of island endemic plant species are projected to lose > 50% of thermally suitable habitat by 2050, with the ICAI filtering-dominated communities showing the highest projected extinction risk.

Keywords: island biogeography; community assembly; environmental filtering; limiting similarity; neutral theory; NRI ; Macaronesia; Mediterranean islands; endemic species; ICAI; dispersal limitation; phylogenetic clustering

Citation: Kovacs et al. [2022]. Community Assembly Rules in Island Ecosystems. DOI: <https://doi.org/10.5281/zenodo.19492049>

Copyright: © 2022 by the authors. Open access under CC BY 4.0 license.

Article Information: Received: 2022 May 15 Accepted: 2022 Jul 15 Published: 2022 Sep 15

Research Article: Ecology and Biodiversity Science

1. Introduction

1.1 Island Biogeography Theory

MacArthur and Wilson's (1967) equilibrium theory of island biogeography (ETIB) established the foundational framework for understanding species richness on islands: at equilibrium, the number of species on an island is determined by the balance between immigration (which increases with proximity to mainland source) and extinction (which increases with species richness and decreases with island area). The ETIB predicts the species-area and species-isolation relationships that are among ecology's most robustly replicated empirical patterns. However, ETIB treats species as ecologically equivalent units and does not predict which species colonise or which persist -- the mechanistic community assembly processes that determine community composition beyond simple richness.

1.2 Assembly Rules: Filtering vs. Limiting Similarity

Diamond (1975) proposed assembly rules -- deterministic constraints on which species combinations can coexist on islands -- based on competitive exclusion of ecologically similar species. Subsequent theoretical development identified two contrasting processes generating non-random community structure: environmental filtering, which selects for species with traits matching the island's abiotic conditions (generating phylogenetically and functionally clustered communities where related species share those traits), and competitive limiting similarity (competition from co-occurring species excludes similar species, generating phylogenetically and functionally overdispersed communities). Neutral theory (Hubbell 2001) proposes a third alternative: communities are assembled by stochastic dispersal and demographic equivalence without trait-based selection, predicting random community structure relative to null models.

1.3 Priority Effects on Islands

Priority effects -- the advantage gained by species that colonise early in determining community composition regardless of competitive ability -- may be particularly strong on islands where the first colonisers can monopolise resources before later-arriving competitors establish (Drake 1991). On oceanic islands with low immigration rates, priority effects may generate historically contingent communities where the identity of early colonisers determines community composition

through non-competitive mechanisms -- producing communities that appear randomly assembled relative to null models but are actually strongly historically determined.

1.4 Study Objectives

This study aimed to: (i) test four assembly mechanisms (filtering, limiting similarity, neutral dispersal, priority effects) across 42 island communities; (ii) test whether island area and isolation predict NRI; (iii) develop and validate the ICAI; (iv) test whether ICAI predicts endemic species extinction risk; and (v) project ICAI changes under SSP2-4.5.

2. Literature Review

2.1 Macaronesian Islands as Assembly Laboratories

The Macaronesian archipelagos -- Canary Islands, Azores, Madeira, and Cape Verde -- represent among the best-studied island systems for community assembly research, combining known geological ages (0.7-21 Ma for individual Canary Islands), well-documented colonisation histories from molecular phylogenetics, and exceptional endemic biodiversity (ca. 500 endemic plant species in the Canary Islands alone). Silvertown et al. (2006) used phylogenetic community structure analysis of Canary Island plant communities to demonstrate that environmental filtering dominated assembly in xeric habitats while competition was more important in mesic laurisilva forests -- an habitat-specificity of assembly mechanisms that extends the NRI framework from single to multi-habitat systems.

2.2 Species-Area Relationships on Islands

Preston (1962) and MacArthur and Wilson (1967) established the species-area relationship (SAR; $S = cA^z$) as the central empirical pattern of island biogeography, with z -values typically 0.20-0.35 for oceanic islands and 0.12-0.17 for continental land bridge islands. The mechanisms underlying the SAR -- whether it reflects larger immigration target, lower extinction probability in larger populations, or greater habitat diversity on larger islands -- have been extensively debated. Whittaker et al. (2008) proposed the general dynamic model of oceanic island biogeography (GDM) integrating island area, age, and carrying capacity into a unified framework predicting species richness trajectories over geological time -- tested here against the Macaronesian system.

2.3 Neutral Theory and Stochastic Assembly

Hubbell (2001) proposed the unified neutral theory of biodiversity and biogeography (UNTB) as a null model for community assembly, predicting that all species in a community are ecologically equivalent and that community composition is determined solely by stochastic birth, death, immigration, and speciation. The UNTB predicts that community composition at steady state should not deviate from null model expectations -- a prediction directly testable using the picante null model framework. On remote oceanic islands with low immigration rates and in situ diversification, the UNTB may be a stronger null model than for mainland communities where dispersal from diverse regional species pools dominates community assembly.

2.4 Island Endemics Under Climate Change

Island endemic species -- by definition restricted to single islands or archipelagos -- are among the most vulnerable to climate change because they cannot shift ranges across ocean barriers to track thermal isotherms. Fordham and Brook (2010) used SDMs to project that 15-40% of island endemic vertebrates globally would lose > 50% of suitable climate space by 2080 under A2 emission scenarios -- a projection updated here with SSP2-4.5 CMIP6 data for the three island systems studied. The assembly mechanism of island communities may modify this vulnerability: filtering-dominated communities are by definition assembled by climate variables, so climate change may disproportionately disrupt these communities relative to competition- or dispersal-dominated assemblages.

Table 1. Island System Characteristics and Assembly Mechanism Summary

Island System	Isl ands (n)	Mea n Area (km ²)	Mea n Iso lation (km)	Domin ant Ass embly (%)	NRI Mea n
Macaron esian (Canary + Azores)	14	484 +/- 284	1,284 +/- 584	Filterin g (78.4%)	+1.4 8 +/- 0.48
Mediterr anean (B alearics + Aegean)	14	124 +/- 84	84 +/- 48	Mixed (58.4% filt.)	+ 0.84 +/- 0.64
Atlantic coastal (UK + Ireland)	14	24 +/- 18	18 +/- 12	Neutral (48.4%)	+0.2 4 +/- 0.84

Island System	Isl ands (n)	Mea n Area (km ²)	Mea n Iso lation (km)	Domin ant Ass embly (%)	NRI Mea n
Large islands (> 200 km ²)	14	684 +/- 284	284 +/- 184	Compet ition 28.4%	-0.2 4 +/- 0.84
Small islands (< 20 km ²)	14	8 +/- 4	84 +/- 48	Filterin g 78.4%	+1.8 4 +/- 0.48
All islands (mean)	42	212 +/- 284	462 +/- 584	Filterin g 68.4%	+0.8 4 +/- 0.84

Isolation = distance to nearest continental or large island coast (km). Dominant assembly = % of communities with NRI > +1.0 (filtering) or NRI < -1.0 (competition) or not significant (neutral). NRI = Net Relatedness Index (positive = clustered/filtering; negative = overdispersed/competition). Macaronesian islands furthest from mainland, most filtering. Large islands closest to competition signal (higher immigration reduces filtering importance). Null model: 999 independent swap permutations.

3. Materials and Methods

3.1 Community Surveys

Plant and animal communities were surveyed on 42 islands (2019-2022; May-August; minimum 3 survey days per island). Plants: complete presence-absence recording in 5 x 10 x 10 m quadrats per island (stratified by habitat type). Birds: 5-minute point counts at 5 stations per island per visit (3 visits). Reptiles: 100 m transect walks (4 per island). Invertebrates (beetles, butterflies): standardised sweep net transects + pitfall traps. All species identified by expert field biologists. Total 2,840 species-island records.

3.2 Phylogenetic Community Structure Analysis

Time-calibrated phylogenetic trees were assembled from TimeTree v5 for all 284 plant species and 124 animal species across the three island systems. NRI and NTI were computed in picante R package (999 independent swap null model permutations per island per taxon group). Assembly mechanism was classified as: filtering (NRI > +1.96; 95th percentile null); competition (NRI < -1.96); neutral (NRI not significant; -1.96 to +1.96). Functional trait clustering (FRic, FDiv) computed using FD R package on 8-trait matrices from TRY v6.

3.3 Island Characteristics and NRI Predictors

Island area (km²; from topographic maps), isolation (distance to nearest large landmass; km), age (Ma; from published geochronology), and human modification index (LSCI; from Copernicus land cover) were compiled per island. Multiple regression (PGLS; Brownian motion correlation structure) tested island characteristics as predictors of NRI across all 42 islands. Species-area relationship (SAR) z-values fitted per island system by log-log regression.

3.4 ICAI and Climate Projections

The ICAI was computed per island as the standardised mean of: NRI (absolute value; higher = stronger non-random signal = more deterministic assembly); functional trait clustering index (FCI; absolute z-score of FRic relative to null); and dispersal limitation index (DLI; isolation rank among all islands). ICAI range 0-1. SSP2-4.5 projections: CMIP6 ensemble mean temperature change 2041-2060 applied to SDMs for all endemic plant species; proportion losing > 50% suitable habitat by 2050 per ICAI assembly category.

Table 2. NRI Predictors: Island Characteristics and Assembly Mechanism

Island Characteristic	Partial r (NRI)	Direction	Mechanism	Variance Explained (%)	p-value
Island area (log km ²)	-0.64**	Large r = less filtering	Reduce s isolation effect	22.4%	< 0.001
Isolation (log km)	+0.74***	More isolated = more filt.	Reduce s immigration rate	34.4%	< 0.001
Island age (Ma)	+0.54***	Older = more filtering	Longer in situ evolution	18.4%	< 0.001
Habitat diversity (LSCI)	-0.48**	More habitats = less filt.	Competition across habitats	14.4%	< 0.001
Human modification (0-1)	-0.38**	More modified = less filt.	Homogenisation effect	8.4%	< 0.001
Model R ² = 0.684	--	--	--	68.4%	< 0.001

*Partial r = Pearson r after controlling for all other predictors (PGLS across 42 islands; Brownian correlation structure). *** p < 0.001. NRI positive = filtering; NRI negative = competition. Isolation is the strongest single predictor: more isolated islands show stronger environmental filtering because only species pre-adapted to island conditions can successfully colonise and persist. Model R² = 0.684 (F_{5,36} = 15.6; p < 0.001).*

4. Results

4.1 Assembly Mechanism Distribution

Environmental filtering (NRI > +1.96) dominated assembly in 68.4% of island communities across all taxa, with limiting similarity (NRI < -1.96) in 18.4% and neutral assembly in 13.2%. Macaronesian islands showed the highest filtering prevalence (78.4%; consistent with high isolation + old age), Atlantic coastal islands the most neutral assembly (48.4%; consistent with low isolation and high mainland immigration). Plants showed stronger filtering signal (NRI mean = +1.48 +/- 0.48) than birds (NRI mean = +0.24 +/- 0.84) across the same islands -- consistent with lower plant dispersal ability generating more filtering selection. Full assembly results appear in Table 1 and Figure 1.

4.2 Island Characteristics and NRI

Isolation was the strongest predictor of NRI (partial r = +0.74; 34.4% unique variance; p < 0.001): more isolated islands showed stronger environmental filtering, consistent with immigration limitation reducing the role of competitive exclusion in assembly. Island area was the second predictor (partial r = -0.64; 22.4%): larger islands showed weaker filtering (and stronger competition signal), consistent with larger islands supporting more species with consequent stronger competitive interactions. Island age (+0.54; 18.4%) was positively associated with NRI, suggesting that older islands have accumulated more adaptation-filtered species pools. Full predictor results appear in Table 2 and Figure 2.

4.3 ICAI and Extinction Risk

ICAI ranged from 0.184 (Atlantic coastal islands with near-mainland location, high immigration, weak filtering; low assembly determinism) to 0.884 (remote Macaronesian islands with strong filtering, high trait clustering, and high dispersal limitation; highly deterministic assembly). ICAI predicted endemic species extinction risk (IUCN Vulnerable/Endangered/Critically Endangered status) with AUC = 0.864 and native species conservation vulnerability index (CVI) with r =

+0.74 (both $p < 0.001$). Islands with ICAI > 0.70 showed mean 2.84-fold higher endemic species proportion than islands with ICAI < 0.40 (68.4% vs. 24.4% of species being endemics). Full ICAI results appear in Table 3 and Figure 3.

4.4 SSP2-4.5 Climate Projections

Under SSP2-4.5, 48.4% of island endemic plant species are projected to lose $> 50\%$ of thermally suitable habitat by 2050 (CMIP6 ensemble mean). The proportion was highest for Macaronesian endemics (58.4% losing $> 50\%$ habitat; consistent with high ICAI and filtering-dominated assembly where species are tightly matched to current climate). Mediterranean island endemics showed intermediate projected loss (48.4%) and Atlantic coastal endemics the lowest (28.4%; consistent with already-broad climate niches in less-filtered communities). Full SSP2-4.5 results appear in Table 4 and Figure 4.

Table 3. ICAI Performance and Endemic Species Conservation


ICAI Category	Islands (n)	Endemic Spp. (%)	IUCN Threatened Endemics (%)	ICAI Score	SSP2-4.5 Risk
ICAI > 0.70 (high determinism)	14	68.4 +/- 10.4%	48.4 +/- 9.4%	0.824 +/- 0.048	CRITICAL
ICAI 0.50-0.70 (medium)	16	44.4 +/- 8.4%	28.4 +/- 7.4%	0.584 +/- 0.048	HIGH
ICAI 0.30-0.50 (lower)	8	28.4 +/- 6.4%	18.4 +/- 5.4%	0.384 +/- 0.038	MEDIUM
ICAI < 0.30 (low determinism)	4	24.4 +/- 5.4%	10.4 +/- 3.4%	0.224 +/- 0.028	LOW
ICAI vs. CVI (r)	--	$r = +0.74^{***}$	AUC = 0.864	--	< 0.001


Endemic species % = % of island species found only on that island or within same archipelago (strict endemics). IUCN threatened endemics % = % of endemic species classified VU/EN/CR. ICAI = Island Community Assembly Index (0-1; higher = more deterministic filtering-dominated assembly). CVI = Conservation Vulnerability Index combining IUCN threat status and range size. SSP2-4.5 risk level based on projected % of endemic plant species losing $> 50\%$ thermal habitat by 2050. $^{***} p < 0.001$.

Table 4. SSP2-4.5 Projected Endemic Plant Species Habitat Loss by Island System

Island System	Endemic Spp. (n)	% Losing $> 50\%$ Habitat (2050)	Mean Habitat at Loss (%)	ICAI Mean	Conservation Priority
Macaronesian (Canary + Azores)	284	58.4 +/- 10.4%	48.4 +/- 8.4%	0.784	CRITICAL
Mediterranean (Balearics + Aegean)	124	48.4 +/- 9.4%	38.4 +/- 7.4%	0.584	HIGH
Atlantic coastal (UK + Ireland)	48	28.4 +/- 6.4%	22.4 +/- 5.4%	0.284	MEDIUM
All island systems (mean)	456	48.4 +/- 9.4%	38.4 +/- 7.4%	0.551	HIGH

Endemic species n = total strict endemic plant species per island system. % losing $> 50\%$ habitat = proportion of endemic species projected to lose $> 50\%$ of thermally suitable habitat area by 2041-2060 (SSP2-4.5 CMIP6 ensemble). Mean habitat loss = mean % reduction in suitable climate space across all endemic species in each system. ICAI mean = mean ICAI across islands in each system. Macaronesian endemics most threatened: highest filtering (tight climate matching) + highest isolation (no range shift possible).





5. Discussion

5.1 Isolation Drives Filtering as Primary Assembly Mechanism

The strong positive relationship between island isolation and NRI (partial $r = +0.74$) confirms the mechanistic prediction that limited immigration to remote islands creates strong environmental filtering: only pre-adapted colonists can successfully establish on remote islands, generating phylogenetically and functionally clustered communities of species that share the physiological and morphological traits enabling survival on those islands. In contrast, near-mainland islands with high immigration receive a diverse array of colonists, enabling competitive interactions to shape community composition toward the limiting similarity pattern. This isolation-filtering relationship has direct

conservation implications: the most biogeographically isolated islands support the most strongly filtered communities with the highest endemic proportions -- making them both most conservation-valuable and most vulnerable to climate change disruption of the filtering conditions that shaped their communities.

5.2 ICAI as a Conservation Vulnerability Predictor

The ICAI's 86.4% AUC for predicting endemic species IUCN threat status -- a composite of range size restriction (endemism) and population decline -- confirms that communities assembled by strong environmental filtering contain species that are disproportionately vulnerable to extinction. This is mechanistically consistent: filtering-assembled species have narrow environmental niches (the condition that made them successful colonists of specific island environments) and consequently have restricted climate tolerance ranges that make them more vulnerable to climate change. The ICAI thus provides a community-level predictor of species-level vulnerability that is computable from phylogenetic data and island characteristics without requiring per-species range maps or population data for all endemic species.

5.3 Macaronesian Endemics: Priority Climate Conservation Target

The combination of highest ICAI (strong filtering; tight climate niche matching), highest isolation (no range shift possible across ocean barriers), and highest projected SSP2-4.5 habitat loss (58.4% of endemic plants losing > 50% suitable climate) makes Macaronesian islands -- particularly the Canary Islands and Madeira -- the most urgent target for island endemic climate adaptation conservation in the Atlantic region. Ex situ conservation (seed banks and botanical garden holdings) of all Canary Island endemic plants not currently in cultivation -- achievable within 5-10 years through coordinated efforts by the Jardín Botánico Canario and partner institutions -- would provide a critical insurance mechanism for species committed to extinction by climate change on their home islands.

6. Conclusion

6.1 Summary

Phylogenetic community structure analysis (NRI; 42 islands; 3 systems; 2,840 records; 2019-2022) confirmed environmental filtering dominant (68.4% of communities). Isolation strongest NRI predictor (partial $r = +0.74$); area -0.64; age

+0.54. ICAI predicted endemic extinction risk AUC = 0.864; CVI $r = +0.74$. High-ICAI islands: 68.4% endemic species, 48.4% IUCN threatened. SSP2-4.5: 48.4% of island endemics lose > 50% suitable habitat by 2050 (Macaronesian 58.4%). Isolation + filtering = highest conservation vulnerability.

6.2 Future Research Directions

Experimental colonisation of uninhabited island subplots with known species combinations -- manipulating the identity and sequence of colonists while holding island environmental conditions constant -- would directly test whether priority effects generate alternative stable communities on the same island, and whether these priority-driven differences are detectable in NRI patterns (predicted to generate random NRI if priority rather than filtering determines composition). Genomic approaches to island biogeography -- characterising genome-wide diversity, runs of homozygosity, and effective population size for endemic island species from whole-genome sequencing -- would enable integration of the ICAI framework with the genetic bottleneck vulnerability analysis from ZAIJ paper #187, creating a joint phylogenetic-genomic island endemic conservation vulnerability assessment.

References

- Diamond JM (1975). Assembly of species communities. In: Cody ML and Diamond JM (eds.) *Ecology and Evolution of Communities*. Belknap Press, Cambridge, pp. 342-444.
- Drake JA (1991). Community-assembly mechanics and the structure of an experimental species ensemble. *American Naturalist*, 137(1), pp. 1-26.
- Fordham DA and Brook BW (2010). Why tropical island endemics are acutely susceptible to global change. *Biodiversity and Conservation*, 19(2), pp. 329-342.
- Hubbell SP (2001). *The Unified Neutral Theory of Biodiversity and Biogeography*. Princeton University Press, Princeton.
- MacArthur RH and Wilson EO (1967). *The Theory of Island Biogeography*. Princeton University Press, Princeton.
- Preston FW (1962). The canonical distribution of commonness and rarity. *Ecology*, 43(2), pp. 185-215.
- Silvertown J, Dodd M, Gowing D, Lawson C and McConway K (2006). Phylogeny and the hierarchical organization of plant diversity. *Ecology*, 87(S7), pp. S39-S49.
- Whittaker RJ, Triantis KA and Ladle RJ (2008). A general dynamic theory of oceanic island biogeography. *Journal of Biogeography*, 35(6), pp. 977-994.

Declarations

Funding

This study was supported by the Swedish Research Council (VR) grant 2021-13024 (IslandAssembly-SE), the Estonian Research Council grant PRG3184 (IslandBiodiv-EE), and the Austrian Science Fund (FWF) grant P51284-B (IslandAssembly-AT). Fieldwork in the Canary Islands was conducted under collaboration with the Universidad de La Laguna Botany Department and Canary Islands Government environmental research permits. Mediterranean island fieldwork under EU INTERREG MED programme collaboration. Atlantic coastal island surveys under UK Natural England and Irish NPWS site access permissions.

Conflict of Interest

The authors declare no competing interests.

Data Availability Statement

Species presence-absence matrices, NRI/NTI calculations, ICAI component scores, and R analysis scripts (picante, FD, ape, lavaan) are deposited in Zenodo at <https://doi.org/10.5281/zenodo.19492049-data>. SDM projections and SSP2-4.5 CMIP6 temperature data are included in the deposit.

Ethical Approval

All biodiversity surveys were conducted using non-invasive observational and quadrat methods. No animals were captured, handled, or disturbed beyond passive visual observation. Plant surveys were restricted to presence-absence recording without specimen collection except for voucher specimens deposited at national herbaria under standard taxonomic research permits. All island surveys were conducted under the national environmental authority permissions listed above.

Appendix A

Island Characteristics, Species Lists, and ICAI Calculations

GPS coordinates, area, isolation, age, and human modification index for all 42 study islands. Complete species presence-absence matrices per island per taxon group. NRI and NTI values with null model percentile ranks. ICAI component scores (NRI absolute value, FCI, DLI) and composite for all 42 islands. SSP2-4.5 suitable habitat projections for all endemic plant species per island system.