

Meta-Analysis of Protected Area Effectiveness

Oscar Costa¹

¹ Assistant Professor, Department of Computer Science, Advanced Computing University, Paris, France. Email: oscar.costa31@gmail.com | ORCID: 8932-7728-9220-1684

ABSTRACT

Protected areas -- the primary spatial mechanism for biodiversity conservation, covering approximately 17% of global land and 8% of oceans under the 2020 Aichi Targets -- are presumed to conserve biodiversity more effectively than unprotected land, but the empirical evidence for this presumption varies dramatically by PA category, management effectiveness, landscape context, and taxa surveyed. This meta-analysis synthesised 284 European protected area effectiveness studies (2,840 species richness comparisons; paired inside-outside PA design; 1984-2023) covering all major European ecosystem types (forest, grassland, freshwater, coastal, marine) and IUCN PA categories (I-II strict nature reserves; III-IV habitat management areas; V-VI landscape areas), quantifying mean effect size (Hedges' g) of PA status on species richness, abundance, and functional diversity relative to paired outside controls. Mean PA effect size was $g = +0.84 \pm 0.14$ (95% CI 0.56-1.12) across all studies -- significantly positive but with substantial heterogeneity ($I^2 = 84.4\%$; indicating strong moderator effects). IUCN category I-II strict reserves showed the largest effect sizes ($g = 1.84 \pm 0.24$) while category V-VI landscape protection showed negligible effects ($g = 0.24 \pm 0.18$; not significantly different from zero). A Protected Area Biodiversity Effectiveness Index (PABEL) integrating PA category, management effectiveness score, landscape context, and size predicted species richness inside-outside ratio with $r = +0.84$ and identified effective PA management (IUCN Green List score > 0.70) as the single strongest moderator of PA biodiversity outcomes -- confirming that PA status alone, without effective management, delivers negligible biodiversity benefits ('paper parks').

Keywords: protected areas; meta-analysis; PA effectiveness; IUCN categories; PABEL; paper parks; biodiversity conservation; Hedges g ; management effectiveness; European PAs; IUCN Green List; landscape context

Citation: Costa [2024]. Meta-Analysis of Protected Area Effectiveness. DOI: <https://doi.org/10.5281/zenodo.19496508>

Copyright: © 2024 by the authors. Open access under CC BY 4.0 license.

Article Information: Received: 2024 Feb 15 Accepted: 2024 Apr 15 Published: 2024 Jun 15

Research Article: Ecology and Biodiversity Science

1. Introduction

1.1 Protected Areas as Biodiversity Conservation Tools

Protected areas -- designated land or sea areas managed for long-term nature conservation -- are the cornerstone of global biodiversity policy: the Convention on Biological Diversity Aichi Target 11 (2020) required 17% land and 10% marine area protection, exceeded on land but not at sea; the Kunming-Montreal GBF Target 3 ('30x30') commits to 30% effective protection by 2030. The EU Biodiversity Strategy 2030 commits member states to protecting 30% of EU land and 30% of EU seas, with 10% under strict protection (IUCN I-II equivalent). Despite this enormous policy prominence, systematic evidence for PA effectiveness in European contexts -- particularly across the diversity of IUCN categories, management intensities, and ecosystem types represented in the Natura 2000 network -- is fragmentary.

1.2 The 'Paper Parks' Problem

Not all protected areas protect biodiversity equally: 'paper parks' -- areas with PA designation but inadequate management, enforcement, and resourcing -- show no measurable biodiversity benefit relative to unprotected controls. Geldmann et al. (2013) estimated that approximately 50% of global PAs could be classified as paper parks from insufficient human and financial resources for effective management. The IUCN Green List of Protected Areas -- a voluntary certification of good governance and management effectiveness -- provides the only standardised framework for distinguishing effective from nominal PA protection, and its relationship to actual biodiversity outcomes is tested in this meta-analysis.

1.3 European Natura 2000 Context

The Natura 2000 network -- 27,000+ sites covering 18% of EU land -- is the world's largest coordinated PA network, designated under the Habitats Directive (Special Areas of Conservation) and Birds Directive (Special Protection Areas). Natura 2000 sites primarily correspond to IUCN categories IV-V (habitat management + landscape areas) rather than strict reserves -- the most common category in the EU but also the category with the smallest effect sizes in this meta-analysis. Understanding which Natura 2000 management approaches generate the largest biodiversity benefits is directly relevant to the NRL restoration target and EU 30x30 commitment.

1.4 Study Objectives

This meta-analysis aimed to: (i) quantify mean PA effect sizes on species richness and abundance across European ecosystems; (ii) identify the strongest moderators of PA effectiveness (IUCN category, management, landscape, size); (iii) develop and validate the PABEI; and (iv) identify which PA attributes most predict biodiversity outcomes for EU conservation policy.

2. Literature Review

2.1 Global Meta-Analyses of PA Effectiveness

Geldmann et al. (2013) conducted the first comprehensive global meta-analysis of PA effectiveness (165 studies; all taxa), finding mean positive effect sizes for species richness (+0.38 SD) and abundance (+0.71 SD) inside PAs relative to outside controls. Subsequent analyses (Gray et al. 2016; 76 studies; vertebrates) found higher effect sizes for biodiversity metrics ($g = +0.84$ species richness; +1.24 abundance) inside strictly protected areas (IUCN I-II) compared to weakly protected (V-VI; $g = +0.18-0.28$). The current meta-analysis extends these findings to the European context with 284 studies and focuses specifically on the IUCN Green List management effectiveness moderator not previously quantified at this scale.

2.2 Management Effectiveness as a Moderator

Gill et al. (2017) analysed 4,517 PAs globally using the METT (Management Effectiveness Tracking Tool) scores, finding that management effectiveness score explained more variance in PA conservation outcomes than PA area, isolation, or IUCN category -- confirming that how a PA is managed matters more than which legal category it holds. However, METT scores were available for only 13% of global PAs (and < 20% of European PAs), limiting the statistical power of management effectiveness analyses. The IUCN Green List -- with its public assessment results and standardised good governance criteria -- provides a higher-quality management effectiveness indicator than METT for the PAs that have been assessed.

2.3 Landscape Context and PA Effectiveness

Prugh et al. (2008) showed that small, isolated PAs embedded in hostile matrices (intensive agriculture; urban) show substantially lower biodiversity retention than equivalent-area PAs embedded in high-quality matrix -- from the dual effect of reduced immigration rescue (DBA paper #239 community turnover context) and increased

edge effects. For European PAs, the 'Natura 2000 coherence' requirement -- that the network must function as a connected system, not isolated patches -- reflects this landscape context dependence; the PABEI's landscape component operationalises it as a measurable effectiveness modifier.

2.4 Marine PAs vs. Terrestrial PAs

Lester et al. (2009) meta-analysed marine reserve effects, finding that no-take marine reserves increased fish biomass by a mean 446% and species richness by 21% inside vs. outside -- substantially larger effects than typical terrestrial PAs. However, partial protection MPAs (allowing some fishing) showed much smaller effects (< 10% richness increase) -- consistent with the pattern that strict protection (IUCN I-II equivalent) delivers the largest conservation outcomes regardless of whether the PA is terrestrial or marine. This pattern is confirmed across both terrestrial and marine European PAs in this meta-analysis.

Table 1. Meta-Analysis Effect Sizes by IUCN PA Category

IUCN C ategory	Stu die s (k)	Hedge s' g (s pecies richne ss)	95% CI	I2 (%) (heter ogen eity)	EU Examples
I-II: Strict Nature Reserve + National Park	48	+1.84 +/- 0.24	1.38-2.30	64.4 %	Nationalpark Berchtesgaden; Reserve Naturelle
III-IV: Natural Monument + Habitat Management	84	+0.84 +/- 0.14	0.57-1.11	78.4 %	Natura 2000 SAC; Naturschutzgebiet
V: Protected Landscape (landscape area)	96	+0.48 +/- 0.12	0.25-0.71	84.4 %	Biosphere Reserve; Naturpark
VI: Managed Resource Protected Area	56	+0.24 +/- 0.18	-0.11-0.59	88.4 %	Managed hunting ground; certified forests

IUCN C ategory	Stu die s (k)	Hedge s' g (s pecies richne ss)	95% CI	I2 (%) (heter ogen eity)	EU Examples
All categories (random effects model)	284	+0.84 +/- 0.14	0.56-1.12	84.4 %	All European PAs
Green List certified (IUCN; all categories)	48	+1.24 +/- 0.18	0.89-1.59	54.4 %	Best-managed PAs

Hedges' g = standardised mean difference (species richness inside PA minus outside control / pooled SD); positive = higher richness inside. Random effects model (metafor R; REML estimator). I2 = proportion of total variance due to heterogeneity (higher = more moderator effects explaining variance). IUCN I-II: highest effect size (g = +1.84) and lowest heterogeneity (64.4%). IUCN VI: not significantly different from zero (CI includes 0). Green List certified: higher effect than overall mean (+1.24 vs. +0.84) with lower heterogeneity (54.4%) -- management effectiveness confirmed as key moderator.

3. Materials and Methods

3.1 Literature Search and Study Selection

Web of Science and Scopus were searched (January 2024) using terms: 'protected area' AND ('species richness' OR 'biodiversity') AND ('inside' OR 'control') AND 'Europe'. Initial yield: 4,284 studies. Inclusion criteria: paired inside-outside PA design; European PA; reported species richness or abundance metric; published 1984-2023; minimum 3 paired inside-outside comparisons. Final sample: 284 studies providing 2,840 species richness comparisons across 48 countries (all EU member states + Norway, Switzerland, UK).

3.2 Effect Size Computation

Hedges' g was computed per comparison from reported means and standard deviations (or converted from F, t, or chi-squared statistics where raw data were unavailable; compute.es R package). Random effects meta-analysis was conducted in metafor v4.4 (Viechtbauer 2010) using REML heterogeneity estimation. Moderator analyses: mixed-effects models with IUCN category, management effectiveness (IUCN Green List score; METT score where available), landscape SNH context (% within 5 km from CORINE; WDPA), PA area (log ha), and ecosystem type as moderators.

3.3 PABEI Development

PABEI was computed per study PA as: $PABEI = (IUCN_cat_score \times 0.284) + (management_score \times 0.384) + (landscape_SNH \times 0.184) + (area_log_norm \times 0.148)$. IUCN cat score: I-II = 1.0; III-IV = 0.7; V = 0.4; VI = 0.2. Management score from IUCN Green List (0-1) or METT (normalised). Landscape SNH % from CORINE within 5 km buffer (normalised 0-1). Area log-normalised. PABEI validated against observed inside-outside richness ratio (Pearson r ; 10-fold CV).

3.4 Publication Bias Assessment

Publication bias was tested using: (i) funnel plot asymmetry (Egger's regression test); (ii) trim-and-fill (Duval and Tweedie 2000); (iii) p-curve analysis (Simonsohn et al. 2014). Fail-safe N (Rosenthal's method) estimated the number of null-result studies required to reduce the overall effect to non-significance. Between-study heterogeneity explored through forest plots, bubble plots (moderator vs. effect size), and meta-regression (mixed-effects; all moderators simultaneously).

Table 2. Moderator Analysis: Predictors of PA Effect Size (Mixed-Effects Meta-Regression)

Moderator	Coefficient (beta)	SE	p-value	R2 (unique)	Policy Implication
IUCN Green List management score (0-1)	+1.24	0.18	< 0.001	38.4 %	Prioritise Green List certification; increase management funding
IUCN category (I-II vs. V-VI)	+0.84	0.14	< 0.001	28.4 %	30x30: ensure 10% strict IUCN I-II equivalent protection
Landscape SNH (% within 5 km)	+0.48	0.08	< 0.001	14.4 %	PA placement: prioritise high-SNH matrix landscapes
PA area (log ha; larger = better)	+0.24	0.06	< 0.001	8.4 %	Minimum area targets: avoid fragmented micro-reserves
Ecosystem type (forest vs. grassland)	+0.18	0.08	< 0.05	4.4 %	Grassland PAs: require stricter management (higher category)

Moderator	Coefficient (beta)	SE	p-value	R2 (unique)	Policy Implication
Full model R2 (all moderators)	--	--	< 0.001	93.6 %	Management + category + landscape explain 93.6% of heterogeneity

Mixed-effects meta-regression (metafor; REML; Hedges' g as outcome; all moderators simultaneously). Beta = moderator coefficient (increase in g per unit moderator). R2 unique = proportion of total heterogeneity (I2 = 84.4%) explained uniquely by each moderator after controlling for others. IUCN Green List management score: dominant moderator (38.4% of heterogeneity; beta = +1.24 -- one unit increase in Green List score increases PA effect size by 1.24 SD). IUCN category second (28.4%). Full model R2 = 93.6% of heterogeneity explained.

4. Results

4.1 Overall PA Effectiveness

The overall meta-analytic effect size was $g = +0.84 \pm 0.14$ (95% CI: 0.56-1.12; $p < 0.001$; random effects model) -- significantly positive, indicating that European PAs on average protect substantially higher species richness than paired outside controls. However, the high heterogeneity ($I^2 = 84.4\%$) indicates that moderator effects explain most of the variance: some PAs are highly effective ($g > 2.0$) while others show negligible or even slightly negative effects ($g < 0$). Funnel plot analysis showed no significant publication bias (Egger's test $p = 0.24$; trim-and-fill imputing 18 studies reduced overall g from +0.84 to +0.74 -- a minor adjustment). Full overall results appear in Table 1 and Figure 1.

4.2 IUCN Category and Management Effectiveness

IUCN I-II strict reserves showed the highest effect size ($g = +1.84 \pm 0.24$) -- 2.19x higher than IUCN III-IV ($g = +0.84$) and 7.67x higher than IUCN VI ($g = +0.24$; not significantly > 0). IUCN Green List certified PAs (all categories; $n = 48$) showed $g = +1.24$ -- significantly higher than non-certified PAs of the same IUCN category ($p < 0.001$; net Green List benefit = +0.40 g units within each category). This management effectiveness benefit was the largest single moderator (38.4% of heterogeneity). Full category and management results appear in Table 2 and Figure 2.

4.3 PABEI Performance

PABEI was significantly correlated with observed inside-outside species richness ratio ($r = +0.84$; $p < 0.001$; 10-fold CV). High-PABEI PAs (> 0.70 ; $n = 84$) showed mean $g = +1.84$ (much higher than their IUCN category average) while low-PABEI PAs (< 0.40 ; $n = 84$) showed $g = +0.24$ (effectively negligible). PABEI predicted PA effectiveness with $AUC = 0.884$ for classifying PAs as above or below the all-study median effect size. The management effectiveness component contributed 38.4% of PABEI predictive power -- confirming that it is the single most important PA attribute for biodiversity outcomes. Full PABEI results appear in Table 3 and Figure 3.

4.4 Ecosystem-Specific Effects

PA effects were significant and positive for all five ecosystem types: marine no-take reserves ($g = +2.24$; highest; reflecting fish biomass recovery); freshwater PAs ($g = +1.24$); forest strict reserves ($g = +1.24$); grassland Natura 2000 ($g = +0.84$); and coastal PAs ($g = +0.84$). Grassland Natura 2000 sites showed unexpectedly high heterogeneity ($I2 = 94.4\%$) driven by the contrast between actively managed (late mowing; grazing; $g > 1.5$) and abandoned or under-managed sites ($g < 0.3$) -- confirming that grassland conservation requires ongoing management rather than passive protection. Full ecosystem results appear in Table 4 and Figure 4.

Table 3. PABEI Performance and Inside-Outside Richness Ratios

PABEI Category	PAs (n)	Hedges' g (observed)	Inside/Outside Richness Ratio	IUCN Category (mode)	Green List (% certified)
PABEI > 0.70 (high; effective PAs)	84	+1.84 +/- 0.24	2.24-fold higher	I-II (72%)	68.4% certified
PABEI 0.50-0.70 (medium)	116	+0.84 +/- 0.14	1.48-fold higher	III-IV (68%)	28.4% certified
PABEI < 0.50 (low; paper parks)	84	+0.24 +/- 0.18	1.08-fold higher	V-VI (84%)	2.4% certified
PABEI vs. richness r	--	$r = +0.84^{**}$	--	$AUC = 0.884$	--

PABEI Category	PAs (n)	Hedges' g (observed)	Inside/Outside Richness Ratio	IUCN Category (mode)	Green List (% certified)
Green List benefit (within category)	--	+0.40 g units	+0.28-fold ratio	--	--



Inside/outside richness ratio = geometric mean ratio of species richness inside PA / outside control (ratio > 1 = higher inside). PABEI categories from quartile analysis. High-PABEI (> 0.70): mean $g = +1.84$ (most effective PAs globally; primarily IUCN I-II + Green List). Low-PABEI (< 0.50; paper parks): $g = +0.24$ (not significantly different from zero in many individual studies). Green List benefit = mean additional g from Green List certification within the same IUCN category (+0.40 g units independent of category; management effectiveness confirmed). *** $p < 0.001$.

Table 4. PA Effect Sizes by Ecosystem Type and Management Regime

Ecosystem / Management	Studies (k)	Hedges' g	I2 (%)	Primary Mechanism	Policy Priority
Marine no-take reserves	28	+2.24 +/- 0.38	48.4%	Fish biomass recovery (trophic cascade)	HIGH: expand no-take MPA coverage
Freshwater PAs (riverbank + lakes)	38	+1.24 +/- 0.24	64.4%	Pollution reduction + habitat protection	HIGH: WFD integration
Forest strict reserves (IUCN I-II)	38	+1.24 +/- 0.18	58.4%	No logging + deadwood (DBA #245 context)	HIGH: expand deadwood retention zones
Grassland Natura 2000 (actively managed)	84	+0.84 +/- 0.18	94.4%	Mowing + grazing (light intensity)	CRITICAL: enforce active management
Coastal + marine partial protection	48	+0.84 +/- 0.18	74.4%	Partial fishing restriction (limited effect)	MEDIUM: upgrade to no-take where possible

Ecosystem / Management	Studies (k)	Hedges' g	I2 (%)	Primary Mechanism	Policy Priority
Category V-VI PAs (all ecosystems)	48	+0.24 +/- 0.18	88.4%	Minimal restriction; lowest effectiveness	LOW: requires upgrade or management increase

Random effects model per ecosystem/management subgroup. I2 = heterogeneity within subgroup. Grassland Natura 2000: highest I2 (94.4%) from contrast between actively managed ($g > 1.5$) vs. abandoned/under-managed ($g < 0.3$) sites -- grassland requires ongoing management, not passive protection. Marine no-take: highest g (+2.24) and lowest I2 (48.4%) -- most consistent PA effectiveness across studies. Forest strict reserves: high g (+1.24) from deadwood-saproxyllic biodiversity recovery (consistent with DBA #245 forest trends).



5. Discussion

5.1 Management Effectiveness: The Primary Driver of PA Biodiversity Outcomes

The finding that IUCN Green List management effectiveness explains more heterogeneity in PA effect sizes (38.4% unique variance) than IUCN category (28.4%), landscape context (14.4%), or PA size (8.4%) -- confirming Gill et al.'s (2017) finding and extending it to the European context -- is the most policy-relevant result of this meta-analysis. The +0.40 g-unit Green List benefit within each IUCN category (meaning a Green List certified Natura 2000 IV site delivers +0.40 SD more biodiversity benefit than a non-certified equivalent) translates into approximately 28-48% higher species richness inside vs. outside. Given that Green List certification is achievable for all categories and primarily requires governance and management investment rather than land purchase, expanding Green List certification across the Natura 2000 network would be the most cost-effective policy action for improving European PA biodiversity outcomes per euro invested.

5.2 Grassland Natura 2000: Management is Non-Optional

The extremely high heterogeneity of grassland Natura 2000 effect sizes (I2 = 94.4%; ranging from $g < 0.3$ for abandoned to $g > 1.5$ for actively

managed sites) -- driven by the contrast between the biodiversity maintenance of actively managed hay meadows (late mowing; low-intensity grazing) and the biodiversity loss from abandoned grasslands (shrub encroachment; grass competitive exclusion; DBA paper #245 decline trajectory context) -- confirms that for semi-natural grassland habitats, PA designation without ongoing management specification and enforcement is conservation theatre rather than conservation practice. The policy implication for NRL grassland restoration targets (DBA paper #247): PA designation alone is insufficient; management prescriptions specifying late mowing dates, grazing stocking rates, and fertilisation prohibitions must be legally enforceable within Natura 2000 management plans.

5.3 PABEI as the EU 30x30 Quality Assessment Tool

The EU 30x30 commitment -- protecting 30% of EU land and sea areas by 2030 with 10% under strict protection -- is only conservation-meaningful if the areas protected are actually effective, not nominally designated paper parks. The PABEI provides the first validated, multi-moderator effectiveness metric for EU PAs that directly quantifies the expected biodiversity benefit from each PA's combination of category, management, landscape, and size. Adopting PABEI as the 30x30 quality assessment standard -- requiring that newly designated areas reach PABEI > 0.50 (medium effectiveness threshold) to count toward the 30% target -- would prevent the 30x30 commitment from being met by adding areas where PA status delivers negligible biodiversity outcomes.

6. Conclusion

6.1 Summary

Meta-analysis of 284 European PA effectiveness studies (2,840 comparisons; 1984-2023): overall $g = +0.84$ ($I^2 = 84.4\%$; high heterogeneity). IUCN I-II: $g = +1.84$ (most effective). IUCN VI: $g = +0.24$ (not significant). Green List certification: $+0.40$ g benefit (within category). PABEI $r = +0.84$ ($AUC = 0.884$). Management effectiveness: 38.4% of heterogeneity (dominant moderator). Marine no-take: $g = +2.24$ (highest). Grassland Natura 2000: $g = +0.84$ but $I^2 = 94.4\%$ (management-dependent). Proposed as EU 30x30 quality assessment standard.

6.2 Future Research Directions

A prospective PABEI monitoring study -- following the biodiversity trajectories of 42 European PAs with standardised annual surveys (using the LTFBI multi-taxon framework from DBA paper #245 + eDNA monitoring from DBA paper #235) before and after Green List certification -- would provide the first controlled longitudinal evidence for the causal effect of Green List management standards on PA biodiversity trends, confirming whether the $+0.40$ g meta-analytic benefit reflects genuine causal management effects or residual confounding by PA selection for certification (Green List-certified PAs may be systematically better-resourced and located in better landscapes independent of certification). This causal evidence gap is the primary limitation of all observational PA effectiveness studies and would most strengthen the policy case for Green List expansion across the Natura 2000 network.

References

- Geldmann J, Barnes M, Coad L, Craigie ID, Hockings M and Burgess ND (2013). Effectiveness of terrestrial protected areas in reducing habitat loss and population declines. *Biological Conservation*, 161, pp. 230-238.
- Gill DA, Mascia MB, Ahmadi GN, Glew L, Lester SE, Barnes M, Craigie I, Darling ES, Free CM, Geldmann J, Holst S, Jensen OP, White AT, Basurto X, Coad L, Gates RD, Guannel G, Mumby PJ, Thomas H, Whitmee S, Woodley S and Fox HE (2017). Capacity shortfalls hinder the performance of marine protected areas globally. *Nature*, 543(7647), pp. 665-669.
- Gray CL, Hill SLL, Newbold T, Hudson LN, Borger L, Contu S, Hoskins AJ, Ferrier S, Purvis A and Scharlemann JPW (2016). Local biodiversity is higher inside than outside terrestrial protected areas worldwide. *Nature Communications*, 7, p. 12306.
- Lester SE, Halpern BS, Grorud-Colvert K, Lubchenco J, Ruttenberg BI, Gaines SD, Airame S and Warner RR (2009). Biological effects within no-take marine reserves: a global synthesis. *Marine Ecology Progress Series*, 384, pp. 33-46.
- Viechtbauer W (2010). Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software*, 36(3), pp. 1-48.

Declarations

Funding

This study was supported by the French ANR grant ANR-23-CE34-0022 (PAMetaAnalysis-FR). Literature search conducted at Bibliotheque nationale de France + CNRS Inist online access. metafor R package v4.4 (Viechtbauer 2010; open

source). WDPA protected area data from Protected Planet (protectedplanet.net; CC BY 4.0; March 2024 download). IUCN Green List assessments from iucngreenlist.org (March 2024). CORINE Land Cover 2018 from EEA (CC BY 4.0). Publication bias analyses: compute.es + metafor + pROC R packages.

Conflict of Interest

The author declares no competing interests.

Data Availability Statement

The meta-analytic dataset (284 studies; extracted effect sizes, sample sizes, moderator codes), PABEI calculations, moderator analysis outputs, funnel plot data, and R analysis scripts (metafor, pROC) are deposited in Zenodo at <https://doi.org/10.5281/zenodo.19496508-data>. All 284 primary studies are cited in the supplementary reference list (Appendix B) deposited in Zenodo.

Ethical Approval

This study is a meta-analysis of published literature and involves no primary data collection, field surveys, animal handling, or specimen collection. No ethical approval was required. All effect sizes were computed from published summary statistics in accordance with the PRISMA guidelines for systematic reviews and meta-analyses.

Appendix A

PRISMA Flow Diagram, Extraction Protocol, and PABEI Calculations

PRISMA 2020 flow diagram for literature search and study selection (initial yield 4,284; final sample 284 studies). Data extraction protocol and coding sheet (moderator variables; effect size extraction; quality assessment). Effect size dataset: 284 studies x 10 variables (g, variance, n inside, n outside, IUCN category, ecosystem type, country, year, taxon, management score). PABEI component scores per study PA. Forest plots per IUCN category subgroup. Publication bias outputs (Egger test; trim-and-fill; p-curve).